

ACKNOWLEDGEMENT

The author sincerely thanks Professor Dr. Keiko T. Natsuaki, Tokyo University of Agriculture, for her encouragement and kind review of this manuscript. Also special thanks to Mr. Komatsubara and Mr. Ali of the Laboratory of Tropical Plant Protection, Tokyo University of Agriculture, Setagaya Campus, for the tremendous help in inputting the DNA data into the DDBJ. The author also thanks Mr. Hiroshi Yoshioka, for providing begonia seedlings for this experiment.

REFERENCES

- De Gruyter, J., A. Auf'm Keller, H. A. Van Kesteren, J. P. Meffert, J. W. Veenbaas-Rijks, and H. N. Cevat. 1994. *Fusarium sacchari* var. *elongatum*—a serious pathogen of Begonia. Verslagen Meded Plantenziektenk Dienst 170: 76 - 79.
- Freeman, S., A. Zveibil, H. Vintal, and M. Maymon. 2002. Isolation of Nonpathogenic Mutants of *Fusarium oxysporum* f.sp. *melonis* for Biological Control of *Fusarium* Wilt in Cucurbits. The American Phytopathological Society. 164 -168.
- Gams, W., E.S. Hoekstra and A. Aptroot. 1998. CBS course of mycology, 4th ed. Centraalbureau voor Schimmelcultures, Baarn, The Netherlands. 1 - 165.
- Gerlach, W., and H. I. Nirenberg. 1982. The genus *Fusarium* – A pictorial atlas. Mitt Biol Bundesanst Land-Forstw Berlin-Dahlem 209:1 - 406.
- Kimura, M. 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequence. J Mol Evol 16: 111-120.
- Leslie, J. F. and B. A. Summerell. 2006. The *Fusarium* Laboratory Manual. Blackwell Publishing.
- Nirenberg, I. H., and K. O'Donnell. 1998. New *Fusarium* species and combinations within the *Gibberella fujikuroi* species complex. Mycologia, 90(3): 434 - 458.
- O'Donnell, K., et al. 2000. Molecular phylogeny of the *Nectria haematococca* – *Fusarium solani* species complex. Mycologia 92: 919 - 938.
- Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mycol Biol Evol 4: 406 - 425.
- Schroers, H. J., K. O'Donnell, R. P. Baayen. et al. 2004. *Fusarium foetens*, a new species pathogenic to begonia elatior hybrids (*Begonia x hiemalis*) and the sister taxon of the *Fusarium oxysporum* species complex. Mycologica 96(2): 393 - 406.
- Sekine, T., H. Kanno, and T. Aoki. 2008 Occurrence of leaf and stem rot caused by *Fusarium foetens* on begonia elatior hybrids (*Begonia x hiemalis*). Japan Journal of Phytopathology.74: 164 - 166.
- Swofford, D. L (1998) PAUP*: phylogenetic analysis using parsimony, Version 4. Sinauer Associates, Sunderland, MA, USA.
- Thompson, J. D., D. G. Higgins, T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Res 22: 46 - 4680.